

Câncer de Endométrio

Carboplatina + Paclitaxel

Carboplatina: AUC 5-7 IV D1
Paclitaxel: 175mg/m² IV D1
a cada 21 dias
Ref. (1)

Adriamicina + Ciclofosfamida

Doxorrubicina: 60mg/m² IV D1
Ciclofosfamida: 500mg/m² IV D1
a cada 21 dias
Ref. (2)

Adriamicina + Cisplatina

Doxorrubicina: 50mg/m² IV D1
Cisplatina: 50mg/m² IV D1
a cada 21 dias
Ref.(3)

Adriamicina + Paclitaxel

Doxorrubicina: 50mg/m² IV D1
Paclitaxel: 150mg/m² IV D1
a cada 21 dias
Ref. (4)

Cisplatina + Doxorrubircina + Paclitaxel

Cisplatina: 50mg/m² IV D1
Doxorrubircina: 45mg/m² IV D1
Paclitaxel: 160mg/m² IV D2
Filgrastima: 5mcg/kg SC D3-D12
a cada 21 dias
Ref. (5)

CAP

Ciclofosfamida: 500mg/m ²	IV	D1
Doxorrubicina: 50mg/m ²	IV	D1
Cisplatina: 500mg/m ²	IV	D1 a cada 21 dias
Ref. (6)		

Carboplatina + Doxorrubicina Lipossomal

Carboplatina: AUC 5	IV	D1
Doxo lipossomal: 40mg/m ²	IV	D1 a cada 28 dias
Ref. (7)		

Doxorrubicina

Doxorrubicina: 60mg/m ²	IV	D1 a cada 21 dias
Ref. (8)		

Paclitaxel

Paclitaxel: 200mg/m ²	IV	D1 a cada 21 dias
Se RXT prévia: 175mg/m ²	IV	D1 a cada 21 dias
Ref. (9)		

Topotecano

Topotecano: 1mg/m ² /dia	IV	D1-D5 a cada 21 dias
Se RXT prévia 0,8mg/m ² /dia	IV	D1-D5 a cada 21 dias
Ref. (10)		

Acetato de Megestrol

Acetato de megestrol: 160mg	VO	Contínuo
Ref.(11)		

Tamoxifeno

Tamoxifeno: 20mg	VO	Contínuo
Ref.(12)		

Anastrozol

Anastrozol: 1 mg/dia VO contínuo

Ref. (13)

Letrozol

Letrozol: 2,5 mg VO dia contínuo

Ref. (14)

Ifosfamida

Ifosfamida: 1200mg/m² IV

Mesna: 300 mg/m² IV D1 a D5 a cada 28 dias
concomitante, 4 e 8hs após ifosfamida

Ref. (15)

Doxorrubicina lipossomal

Doxorrubicina lipossomal: 40mg/m² IV D1 28 dias

Ref. (16)

1. Hoskins, P.J., et al. Paclitaxel and carboplatin, alone or with irradiation, in advanced or recurrent endometrial cancer: a phase II study. *J Clin Oncol*, 2001. 19(20): p. 4048-53.
2. Thigpen, J.T., et al. A randomized comparison of doxorubicin alone versus doxorubicin plus cyclophosphamide in the management of advanced or recurrent endometrial carcinoma: A Gynecologic Oncology Group study. *J Clin Oncol*, 1994. 12(7): p. 1408-14.
3. Deppe, G., et al. Treatment of recurrent and metastatic endometrial carcinoma with cisplatin and doxorubicin. *Eur J Gynaecol Oncol*, 1994. 15(4): p. 263-6.
4. Fleming G. F., Filiaci V. L., Bentley R. C, et. al. Phase III randomized trial of doxorubicin + cisplatin versus doxorubicin + 24-h paclitaxel + filgrastim in endometrial carcinoma: a

- Gynecologic Oncology Group study. Ann Oncol. 2004; 15(8): 1173-1178.
- 5.** Fleming, G.F., et al. Phase III trial of doxorubicin plus cisplatin with or without paclitaxel plus filgrastim in advanced endometrial carcinoma: a Gynecologic Oncology Group Study. J Clin Oncol, 2004. 22(11): p. 2159-66.
 - 6.** Burke, T.W., et al. Postoperative adjuvant cisplatin, doxorubicin, and cyclophosphamide (PAC) chemotherapy in women with high-risk endometrial carcinoma. Gynecol Oncol, 1994. 55(1): p. 47-50.
 - 7.** Pignata, S., et al. A multicentre phase II study of carboplatin plus pegylated liposomal doxorubicin as first-line chemotherapy for patients with advanced or recurrent endometrial carcinoma: the END-1 study of the MITO (Multicentre Italian Trials in Ovarian Cancer and Gynecologic Malignancies) group. Br J Cancer, 2007. 96(11): p. 1639-43.
 - 8.** Muss, H.B., Chemotherapy of metastatic endometrial cancer. Semin Oncol, 1994. 21(1): p. 107-13.
 - 9.** Ball, H.G., Do we know the best therapy for early endometrial cancer? Gynecol Oncol, 1996. 60(2): p. 173-5.
 - 10.** Wadler, S., et al. Topotecan is an active agent in the first-line treatment of metastatic or recurrent endometrial carcinoma: Eastern Cooperative Oncology Group Study E3E93. J Clin Oncol, 2003. 21(11): p. 2110-4.
 - 11.** Thigpen, J.T., et al. Oral medroxyprogesterone acetate in the treatment of advanced or recurrent endometrial carcinoma: a dose-response study by the Gynecologic Oncology Group. J Clin Oncol, 1999. 17(6): p. 1736-44.
 - 12.** Thigpen, T., et al. Tamoxifen in the treatment of advanced or recurrent endometrial carcinoma: a Gynecologic Oncology Group study. J Clin Oncol, 2001. 19(2): p. 364-7.
 - 13.** Rose PG, Brunetto, VL, et al. A Phase II trial of Anastrozole in advanced recurrent or persistent endometrial carcinoma: A Gynecologic Oncology Group Study. Gynecologic Oncology, 2000; 78, 212-216.

- 14.** Ma B B.Y, Oza A, *et al.* The activity of letrozole in patients with advanced or recurrent endometrial cancer and correlation with biological markers – a study of the National Cancer Institute of Canada Clinical Trials Group, Intern J Gynecol Cancer 2004; 14(4):650 - 658.
- 15.** Sutton GP, *et al.* A Phase II Gynecologic Oncology Group Trial of Ifosfamide and Mesna in Advanced or Recurrent Adenocarcinoma of the Endometrium. Gynecol Oncol. Ginecol Oncol 1996;63:25-7.
- 16.** Homesley, HD, *et. al.* Phase II trial of liposomal doxorubicin at 40 mg/m² every 4 weeks in endometrial carcinoma: A Gynecologic Oncology Group Study Gynecol Oncol 2005; 98:294-8.